

TARGATT™ CHO-K1 Knock-in Kit (H11) for Antibody Library Screening (Drug Selection)

The TARGATT™ CHO-K1 Kit (H11) for Antibody Library Screening (Drug Selection) was designed for fast and site-specific knock-in in Chinese Hamster Ovary (CHO) cells using an easy-to-use gene knock-in approach. The master cell line provided in this kit contains the attP docking-site engineered into the CHO H11 safe harbor locus in the genome. Any gene of interest can be cloned into the provided TARGATT™ attB cloning plasmid, and transfected into the master cell line for generating a stable knock-in cell line. The TARGATT™ integrase-based technology enables efficient DNA integration and high-level gene expression without disrupting internal genes. The TARGATT™ CHO-K1 cell line can therefore be used for uniform, site-specific gene knock-in, generation of isogenic cell lines, and amplification strategies for isolating high expression cells, without the need for single cell cloning.

Advantages of using TARGATT™ Master Cell Lines for gene knock-in:

- High efficiency, unidirectional integration
- Site-specific, stable knock-in cell line generation
- Single copy gene integration into safe harbor locus
- Gene expression from an active, intergenic locus
- Easy-to-use protocol requiring transfection of only one (donor) plasmid

The TARGATT™ CHO-K1 Kit (H11) for Antibody Library Screening (Drug Selection) includes the TARGATT™ 40 attB-Blas-P2A-LacZ (library) Cloning Plasmid (AST3080) which contains blasticidinR. This kit is ideal for drug selection and is suitable for research applications involving gene library construction.

** TARGATT™ master cell lines can be generated in any cell line including stem cells. Please contact Applied StemCell for TARGATT™ cell line engineering services to generate a master cell line in a specific cell line of choice.*

**The TARGATT™ CHO-K1 Kit (H11) for Antibody Library Screening (FACS) is also available. The cloning plasmid in this kit contains mCherry and is ideal for fluorescence-activated cell sorting (FACS).*

Parental Line CHO-K1 Suspension Cells

Catalog #	Product	Size
AST-1410	TARGATT CHO-K1 Knock-in Kit (H11) for Antibody Library Screening (Drug Selection)	1 kit

Intended Use

This product is for research use only and not intended for human or animal diagnostic or therapeutic uses.

Quality Assurance

A certificate of analysis (COA) with detailed quality control information for the cell line and components of the kit will be provided with each shipment.

Storage Instructions and Stability

Store the TARGATT™ master cell line in liquid nitrogen freezer immediately upon receipt. Store the plasmids at -20°C. Do not freeze-thaw plasmids repeatedly. The products provided in this kit are stable for at least 6 months from the date of receiving when stored as directed.

Precautions

PLEASE READ BEFORE HANDLING ANY FROZEN VIALS. Please wear appropriate Personal Protection Equipment (lab coat, thermal gloves, safety goggles and a face shield) when handling frozen vials. Please be aware that the following scenario can occur: Liquid nitrogen can leak into the vials when the vials are submerged in liquid nitrogen. Upon thawing, the liquid nitrogen returns to the gas phase, resulting in a dangerous build-up of pressure within the vial. This can result in the vial exploding and expelling not only the vial contents but also the vial cap and plastic fragments of the vial.

Directions for Use

Kit Components

Catalog #	Component	Amount
AST-1408-1	TARGATT™ CHO-K1 (H11) Master Cell Line	1 x 10 ⁶ cells
AST-3203	TARGATT™ CAG Integrase Plasmid ^{#†}	60 µg
AST-3080	TARGATT™ 40 attB-Blas-P2A-LacZ (library) Cloning Plasmid [#]	15 µg
AST-3065	TARGATT™ 25 Positive Control Plasmid ^{#†}	15 µg

Aliquot into single-use quantities and store at -20°C; do not freeze-thaw the plasmid repeatedly. † The positive control and integrase plasmids are proprietary materials and should not be amplified. Additional quantities of these plasmids are available for purchase.

1. Preparation of Medium

CHO Culture Medium (Complete Culture Medium)

- BalanCD® CHO Growth A Medium
- L-Glutamine (8 mM)

Cryopreservation/ Freezing Medium

- CHO Culture Medium (90%)
- DMSO (10%)

2. Thawing and Culturing Cryopreserved CHO-K1 Cells

- 2.1. Aliquot 30 mL CHO culture medium into a 50 mL conical tube, and warm in a 37°C water bath for 10 minutes. Before transferring into the hood, wipe the tube with paper towel and again with paper towel sprayed with 70% ethanol to disinfect.
- 2.2. Retrieve a vial of frozen TARGATT™ CHO-K1 master cell line from the liquid nitrogen tank

- wearing a thermal glove.
- 2.3. Immediately immerse the vial in a 37°C water bath. While holding the top of the vial, gently agitate the vial.
Note: Gently agitate the vial to thaw the cells quickly; DO NOT keep it static while in the water bath; DO NOT submerge the cap.
 - 2.4. When almost completely thawed (usually takes ~1 minute, a small piece of ice is still visible), remove the vial from the water bath, and decontaminate the exterior of the vial with 70% ethanol.
Note: All further operations should be carried out under aseptic conditions
 - 2.5. Transfer the contents of the vial (1 mL of cell suspension) to a conical tube containing 4.0 mL of pre-warmed complete culture medium.
 - 2.6. Determine the viable cell count and calculate cell density, using an appropriate cell counter.
 - 2.7. Centrifuge at 125 x g for 5 minutes. Discard the supernatant.
 - 2.8. Tap the tube to loosen up the cells and re-suspend the cell pellet in pre-warmed fresh complete culture medium at a density of $2.5 - 3.0 \times 10^5$ viable cells/mL, in a fresh 125 mL Erlenmeyer shaker flask.
 - 2.9. Place flask on an orbital shaker platform rotating at 130 rpm, in a 37°C and 5-8% CO₂ incubator.

3. Passaging Procedure

- 3.1. Once the culture reaches $2.5 - 5.0 \times 10^6$ cell/mL, the cells are ready to be passaged.
- 3.2. Warm-up complete culture medium in a 37°C water bath for 10 minutes.
- 3.3. Take the 125 mL Erlenmeyer shaker flask out of the CO₂ incubator.
- 3.4. Determine the viable and total cell counts.
- 3.5. Seed the cells at a density of $2.0 - 3.0 \times 10^5$ viable cells/mL into pre-warmed fresh culture medium by gently pipetting.
Note: Calculate the volume of cell culture suspension and fresh complete medium needed to seed each new flask by dilution.
- 3.6. Passage cells every 3-4 days.
- 3.7. Incubate cultures at 37°C and 5-8% CO₂.

4. Cryopreserving cells

- 4.1. Harvest cells and transfer the contents of the vial into a centrifuge tube.
- 4.2. Centrifuge at 125 x g for 5 minutes.
- 4.3. Aspirate the supernatant and resuspend the cells in cryopreservation medium (refer to step 1).
- 4.4. Aliquot 1 mL of the cell suspension into cryovials and transfer the vials into a CoolCell® cell freezing container and store the vials into a -80°C freezer overnight.
- 4.5. Ensure cells are completely frozen and transfer the cryovials into liquid nitrogen for long-term storage.

5. Transfection: Fast Knock-in Procedure Using Neon™ Transfection System

Cloning with the TARGATT™ 40 attB-Blas-P2A-LacZ (library) Cloning Plasmid

- 5.1. Clone the gene of interest into the TARGATT™ 40 attB-Blas-P2A-LacZ (library) Cloning Plasmid to make the donor plasmid using the NEB® Golden Gate Assembly Kit (Bsal-HF®v2).
- 5.2. Amplify the plasmids using E. cloni® 10G SUPREME SOLOs electrocompetent cells
Note 1: See the Lucigen E. cloni® 10G and 10GF' Electrocompetent Cells manual for the transformation protocol.
Note 2: Do not use 5-alpha competent cells to transform the plasmids.
- 5.3. Purify plasmid pool, aliquot into single-use quantity, and store at -20°C. Do not freeze-thaw the plasmid repeatedly

Preparation of Neon™ Transfection System

- 5.4. Aliquot a sufficient amount of Buffer E2 and Buffer R and warm at room temperature for at least 30 minutes.
- 5.5. Turn on the Neon™ Transfection System 30 minutes before use.

Cell Preparation for Electroporation

- 5.6. Pre-warm 6-well plates with 3 mL of fresh cell culture medium per well at 37°C.
- 5.7. Using cells already in the culture medium, count the cells and ensure >90% cell viability.
- 5.8. Harvest 1.1×10^6 cells and centrifuge the cells at 125 x g for 5 minutes.
- 5.9. Aspirate the supernatant and wash the cells by resuspending in 5 mL PBS.
- 5.10. Centrifuge the cell suspension at 125 x g for 5 minutes. Aspirate the supernatant.
- 5.11. Resuspend the cells by adding 110 µL Neon® Buffer R.
- 5.12. Aliquot a 110 µL cell suspension into a 1.5 mL Eppendorf tube and add 7.5 µg cloned TARGATT™ donor plasmid containing your gene of interest and 2.5 µg TARGATT™ CAG integrase plasmid).
- 5.13. Flick the tube a few times to mix them well.
- 5.14. Electroporate the cells based on the following condition: 1600 V, 30 ms, 1 Pulse with Neon™ electroporation (or any other transfection method-of-choice).
Note: If using other transfection methods, please optimize your protocol accordingly.
- 5.15. After electroporation, plate the cells in one-well of 6-well-plate, and transfer the plate into an incubator at 37°C with 5-8% CO₂. More than one cell well can be done per round of electroporation to increase the library size.
- 5.16. At day 2-3 after electroporation, discard the spent medium and add 3 mL fresh, complete culture medium per well.
- 5.17. Place the culture plate containing the cells into an incubator at 37°C with 5-8% CO₂, on an orbital shaker platform rotating at 130 rpm.

6. Enrichment Using Blasticidin Selection

- 6.1. Seven (7) to 10 days after transfection when cell viability is above 90%, adjust the cell concentration to 1×10^6 cells/mL in 3 mL of culture medium in a 6-well plate.
- 6.2. Add 10 µg/mL of Blasticidin in the medium.
- 6.3. On day 3 of selection, harvest the cells by centrifuge at 125 x g for 5 minutes.
- 6.4. Remove the supernatant and resuspend cells in fresh culture medium with 10 µg/mL of Blasticidin.
- 6.5. Subculture the cells at a cell density of 1×10^6 cells/mL into a 6-well culture plate.
- 6.6. Monitor the cell viability daily during Blasticidin selection.
- 6.7. Keep Blasticidin in the medium for 5-7 days until cell viability is about 30% and then remove Blasticidin from the culture medium.
- 6.8. To stop the Blasticidin selection, harvest the cells and centrifuge the cell suspension at 125 x g for 5 minutes.
- 6.9. Remove the supernatant and resuspend the cells in fresh culture medium, and culture the cells in a 6-well plate until cells recover fully.
- 6.10. Passage the cells when the cell density is above 5×10^6 cells/mL.




Figure 1. Schematic representation of TARGATT™ site-specific transgene integration mediated by integrase in the TARGATT™ CHO-K1 Master Cell Line. The TARGATT™ CHO-K1 (H11) Master Cell Line was engineered with the attP landing pad at the CHO H11 safe harbor locus. The TARGATT™ donor vector containing the attB site is used to clone the transgene. The integrase from the TARGATT™ Integrase Plasmid catalyzes an irreversible reaction between the attP site in the genome and attB site in the donor vector, resulting in site specific integration of the gene of interest at the selected locus.




Figure 2. Genotyping of the 5'-arm and 3'-arm in TARGATT™ CHO-K1 master cell line. PCR analysis shows the expected sizes for the 5' arm and 3' arm junction PCR product in a 1.5% agarose gel: 662bp and 915bp, respectively. Primer sequences used: 5'-arm-Forward: 5'-GGTTGATGACCCTGCACATAGC-3'; 5'-arm-Reverse: 5'TGGATTTTGACTGCAGGGGTAAA-3'; 3'-arm-Forward: 5'-AGGTGTCTGCAGGCTCAAAGAG-3'; 3'-arm-Reverse: 5'AGGCTGGAGGTAGTTTGTG-3'.




Figure 3. Flow cytometric analysis of mCherry expression level in CHO-K1 cells. A donor plasmid containing the fluorescent marker mCherry was used to analyze integration efficiency. An integration efficiency of ~18% without drug selection (right) and a negative control integration efficiency of ~1.60% (left) were obtained.

Related Products

Catalog #	Product	Size
AST-1409	TARGATT CHO-K1 Knock-in Kit (H11) for Antibody Library Screening (FACS)	1 kit
AST-1405	TARGATT CHO-K1 Knock-in Kit (A2)	1 kit
AST-1420	TARGATT CHO-K1 Knock-in Kit (H11)	1 kit

Technical Support

CONTACT US

For more information or assistance, contact Customer Service at:

- Email: info@appliedstemcell.com
- Direct line: [\(408\) 773-8007](tel:4087738007)

WEBSITE RESOURCES

Visit appliedstemcell.com for technical resources and information including:

- Safety Data Sheets (SDS)
- Certificates of Analysis (when available)
- FAQs
- Product literature
- Complete contact information

Applied StemCell, Inc.

521 Cottonwood Dr, Suite # 111, Milpitas, California, 95035, USA • Telephone: +1 866.497.4180 • appliedstemcell.com

© 2026 Applied StemCell, Inc. All rights reserved. TARGATT™ and ActiCells™ are trademarks of Applied StemCell, Inc. All other trademarks are the property of their respective owners. PN: IFU-AST-1410 Rev. 1

Effective Date: 15 September 2026